Content available at: <https://www.ipinnovative.com/open-access-journals>

Indian Journal of Obstetrics and Gynecology Research

Journal homepage: www.ijogr.org

Original Research Article

Sexual practices in pregnant women belonging to low socio-economic status at a tertiary care hospital in New Delhi, India

Chanda Rai^{1,*}, Ananya Banerjee¹, Manisha Meena¹¹Dept. of Obstetrics & Gynecology, VMMC & Safdarjung Hospital, New Delhi, India

ARTICLE INFO

Article history:

Received 03-11-2020

Accepted 27-11-2020

Available online 13-03-2021

Keywords:

Sex in pregnancy

Coitus in married couples

Sexual health

ABSTRACT

Introduction: Sexual changes in pregnancy remains a rarely discussed topic here in India because of social taboos and the embarrassment couples face in discussing these issues.**Aim:** To highlight the sexual changes that take place in all three trimesters of pregnancy and to evaluate the various factors which influence it.**Materials and Methods:** A cross-sectional descriptive study was carried out in the department of Obstetrics & Gynecology at VMMC & Safdarjung Hospital, New Delhi. Two hundred and five consenting women were asked open end questions about their sexual health and variation in the different trimesters. Women with complications like any bleeding or leaking in pregnancy were excluded from the study. Factors affecting their sexual life were taken into account. Data was summarized using descriptive statistics and analysed in SPSS software.**Results:** Our study showed a decline in the frequency of sexual activity by the third trimester with the percentage of women engaging in sexual activities in the first, second and third trimester being 36.6%, 43.9% and 6.3% respectively. Around 9.8% women totally refrained from coitus throughout pregnancy. Libido was found to decrease in 38% of women. The prohibiting factors for coitus included the risk of preterm labour, bleeding and chances of infection to the baby. None of these couple thought of seeking a health professional's advice for their sexual problems.**Conclusion:** Counselling for sexual health must be initiated by the health workers as it will help couples who are embarrassed to discuss these issues owing to the social and cultural norms. This will help to bridge the communication barrier and will be instrumental to a healthy sexual life of married couples who are expecting a child.**Key Messages:** Sexual health in pregnant women largely remains a less discussed topic.

There is a decline in coital frequency during pregnancy.

There are many factors like fear of harming of the fetus which women do not discuss with their health-care providers.

© This is an open access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>) which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

1. Introduction

Sex in pregnancy remains a rarely discussed issue majorly because of the societal norms and embarrassment in asking queries relevant to sex. A pregnant woman undergoes changes both physically and psychologically because of the hormonal changes which is seen to affect her sexual life.^{1,2} The biological changes in pregnancy brought by the

hormonal surge of estrogen, progesterone and prolactin is known to decrease their libido.^{1,3,4}

In our country discussing about sex and sexual health is considered a taboo, thus preventing couples from coming forward and discussing sex related problems with their physicians. Even physicians shy away from this topic. This study aims to bring forth the hidden problems involving the sexual life of a married couple who are eagerly waiting for parenthood. Sex in pregnancy is feared by a lot of couples mainly because of the fear of abortion or preterm labour

* Corresponding author.

E-mail address: chanda17.rims@gmail.com (C. Rai).

or other pregnancy complications.^{5–7} Maintaining sexual relations in pregnancy also depends on a very large extent, influenced by the commands of elders at home, especially in countries like India where societal and religious beliefs play a very important role in decision making. These myths and notions involving sex in pregnancy needs to be dissolved so as to ensure a healthy and blissful sexual life of a couple.

Many pregnant woman are uncomfortable and shy in discussing sexual health during pregnancy with the health workers and doctors and for many pregnancy is a time for no-sex because of multiple reasons. This is not confined to Indian women but other Asian countries like China also mention about the embarrassment faced by their women when they had to confront questions about sexual health and the various reasons which prohibit sexual activities in pregnancy.^{2,8–10}

This study is an attempt to bring to the forefront the sexual changes which take place in all three trimesters of pregnancy and to discuss the various factors which influence sexual activity in a cross-sectional population of Indian women where the societal and cultural beliefs have a huge impact on their personal beliefs and behaviour. There is limited knowledge about sexual and reproductive health in Indian women of reproductive age group particularly in rural India where 92% of the married women had no say in their general health care and 94% could not choose their own doctor.¹¹ Such a serious plight of health awareness in India calls for a unified effort to bring to the notice of all health care professionals various facts about the sexual health in pregnant woman which will further help them in providing proper advise and counselling related to sexual health.

This study is an attempt to understand the attitude, knowledge and behaviour of pregnant women regarding their sexual life, the problems they face, the fears they encounter, what holds them from carrying on with their normal sexual behaviour and ways to dispel these psychological issues. Thus, we have attempted to bridge the gap between expectant mothers and the health care provider concerning their sexual well-being.

2. Materials and Methods

We did a cross-sectional observational study in the Obstetrics and Gynecology department at VMMC and Safdarjung Hospital, New Delhi over a period of thirteen months from March 2016 to August 2017. This hospital is one of the largest tertiary care hospital in New Delhi with over 25,000 deliveries a year and being a government funded institute caters mainly to the poor patients.

The representative sample consisted of 205 pregnant woman who came to the hospital OPD for regular check-up and inpatients who were admitted for various reasons. All consenting gravid women in different trimesters of pregnancy without any pregnancy complications were

included in the study. Women with complications like early pregnancy bleeding, ante-partum haemorrhage, leaking per vaginam were excluded from this study. Women were asked open end questions about their sexual habits in pregnancy and the various factors which affected it. Information like age, gravida, parity, education, socio-economic status, previous obstetric history, the frequency of sex, sexual interest and factors which influence their sexual behaviour could be elicited from the questionnaire. They were also enquired about their knowledge of sexual health and any opinion sought about by their health care professionals. The primary aim of this study was to find out about the sexual practises in women in all three trimesters and to evaluate the different factors which have an impact on sexual behaviour.

Categorical variables are shown using percentage. Graphical representation of nominal data was done using bar graphs. Numerical data was shown using median (IQR). Student independent t -test was done to test the association of numerical data. The level of statistical significance was $p < 0.05$. Outcome variables included frequency of having sex two or less than two times during pregnancy and more than three times during pregnancy. Data was analysed in SPSS version 25.

3. Results

The socio-demographic profile of the patients in their third trimester has been listed in Table 1.

Table 1: Socio-demographic details

	Median	IQR	F (df)	p value
Age in years	24 year	6	0.25(1,203)	0.618
Gravida	2	2	0.75(1,203)	0.430
Parity	0	1	4.5((1,203)	0.015
Per capita income (₹)	11000	5500	1.75(1,203)	0.192

3.1. Sexual activity in pregnancy

Table 2 summarises the coital frequency throughout the duration of pregnancy. Out of the 205 participants, eleven women (5.4%) never had any sexual activity throughout pregnancy. Around 101 (49.2%) of them, had coitus one to two times and the remaining 93 participants (45.4%) could have sexual intercourse for more than three times during the entire period of pregnancy.

Table 2: Frequency of sexual Intercourse during pregnancy

Frequency of Coitus	N	Percentage
No coitus	11	5.4
1-3	101	49.2
>3	93	45.4

We observed a reduction in the frequency of sexual activity in the third trimester with only 13/205 women (6.3%) reporting it, and another 75/205 women (36.6%) and 90/205 (43.9%) could continue sexual practices in the first and second trimesters, respectively. For 16/205 participants, sex was possible in all three trimesters (Table 3). It was seen that around 10% women had gone to their mother's house on confirmation of pregnancy for safe confinement which is a common practice in India and especially in cases of first pregnancy.

Table 3: Trimester wise coital frequency

Trimester	N	Percentage
First	75	36.6
Second	90	43.9
Third	13	6.3
All three trimester	16	7.8
No coitus	11	5.4

3.2. Types of sexual activities

The various types of sexual practices are encapsulated in Table 4. It was observed that majority of the study participants (75.6%) indulged in vaginal intercourse. Among the study population, 25/205 (12.2%) of the women indulged in oral sex and 17/205 (8.3%) found sexual appeasement in masturbation. It was also revealed that 8/205 (3.9%) of the women practised anal intercourse.

Table 4: Type of sexual activity

Type of Sexual Activity	Number	Percentage
Vaginal Intercourse	155	75.6
Oral	25	12.2
Masturbation	17	8.3
Anal	8	3.9

Libido, the desire to have coitus, during pregnancy was found to decrease in majority of the participants while it remained the same as prior to conception in 63/205 (30.7%) of the women. Sexual urge was found to increase in 64/205 (31.3%) of women (Figure 1)

3.3. Prohibiting factors for sex

When we tried to reason out the various causes which caused the inhibition in sexual activities during pregnancy we found that the risk of onset of preterm labour, bleeding and the chances of infection to the baby remained the most cited ground for non-indulgence in sexual activity seen in 42/205, 74/205, 44/205 women respectively. Another 18.5% quoted the fear of injuring the fetus as a factor for hindering sexual activities (Table 4). Only seven (3.4%) study participants had no problem in practicing sex.

Fig. 1: Effect on Libido during pregnancy

Table 5: Factors prohibiting sexual activity during pregnancy

Concerns	Number	Percentage
Injury to Foetus	38	18.5
Early Onset of Labour	42	20.5
Infection	44	21.5
Bleeding	74	36.1
None	7	3.4

3.4. Perineal hygiene practises

Figure 2 reviews the perineal hygiene practices commonly carried out. Another revelation was that many patients used soap and water (32.6%) and antiseptic mostly. Dettol and water (3.9%) for their perineal hygiene. More than half of the population used water only for cleaning their perineum. These women had no formal knowledge on the correct practices for maintaining vaginal hygiene.

3.5. Sources of information

When we enquired about their source of information about sex in pregnancy, more than half of women relied on the advice of the elders in their family and only 25% sought the help of a health-care professional. Majority of them did not seek any medical advice and were dependent on the commands of their family members. (Figure 3)

4. Discussion

The study shows a definite decline in the coital frequency towards the third trimester of pregnancy which has also been seen in other studies.^{5,12–14} This can be attributed to

Fig. 2: Perineal hygiene practices in women

Fig. 3: Source of information

the reduction in libido in many females during pregnancy which was also seen in other studies.^{5,12,15} Erol et al. found diminished clitoral sensation as the most common sexual dysfunction followed by lack of libido and orgasmic disorder. He found no correlation between female sexual function scores and serum androgen levels in pregnancy.¹⁶

Systematic reviews by Von Sydow¹⁷ and Serati et al.¹⁸ have shown that the frequency of coitus does not change or changes only minimally during the first and second trimesters of pregnancy. A study on Taiwanese women has shown that 13.9% of women ceased coitus completely as compared to 9.8% in our study.¹⁹

Many studies have also shown that the both the frequency and the desire for sexual intercourse reduced in third trimester.^{6,7,20} This depends on many physical factors and

hormonal changes of estrogen, progesterone and prolactin which causes breast tenderness, weight gain and fatigue which in turn causes difficulty in getting aroused.^{5,13,21}

Many women also gave history of non-consensual sex in our study. Similar study showed pressured sex rate of 19%.⁵

A look into the causes deterring the carnal desire in women is the fear of harming the baby or the risk of early setting in of labour pain or risk of infection or bleeding. Many studies have highlighted decrease in sexual activity in couples because of the fear of pregnancy complications in both the partners.^{5-7,15}

Proper counselling and approach by the doctors regarding these issues will help women discuss their physical as well as psychosexual problems, concerns regarding decrease in sexual activities and any other related problems. Indian women are particularly shy and feel embarrassed to discuss sexual problems. An open free relationship will help to bridge this gap of communication. This type of embarrassment in discussing their sexual life is also found in women of other Asian countries like Chinese women who feel shy to discuss topics related to their sexual life with their doctors.^{9,10}

Then there is lack of awareness and proper information about clean vaginal practices. Effort must be made to discuss these issues.

Vaginal intercourse remained the most common type of sexual intercourse as seen in other studies.⁵ Some studies noted a decrease in vaginal intercourse and rise in other forms of sexual activity like manual sex, anal and oral sex because of the fear of harming the fetus.²

Many women seek safe confinement in pregnancy because of which they often go back to their mother's family, as seen in our study. This was also seen in Chinese women where sexual prohibitions during pregnancy are aimed to avoid problems in pregnancy.⁸ Many mothers and mother in laws play a pivotal role in decision making for the couple in India thus indicating that social and cultural issues cannot be overlooked, as was also pointed by Pautela et al.¹ In other Asian countries like China and Taiwan, too familial influence seems to play a significant role in decision making.^{9,10}

5. Limitations

The present is an observational study carried out among women belonging to low socio-economic status visiting only one tertiary care hospital, because of this small sample size the study cannot be generalized to the entire cohort of pregnant women belonging to different strata of society.

6. Recommendation

The sexual life of a married couple, especially in India, remains largely an untouched topic. Couples are reluctant

to come forth with their problems owing to societal norms and personal embarrassment. Health-care professionals, too, have not been trained to counsel such couples and create an atmosphere congenial to discuss such issues and not to brush them aside. It is imperative that medical professionals come forward to help these couples discuss their sexual lives freely and devote some time to ease their married lives.

7. Conclusion

A healthy approach for the patients to discuss all sorts of sexual and reproductive health problems should be inculcated. The initiative can be taken from the side of the doctors to promote a uninhibited discussion on these issues to newly married couples and to all couples in general in order to allay their unspoken fears regarding their sexual relationship and the impact on their baby. The emotional and physical aspects of the patients should be dealt with simultaneously to encourage a healthy sexual life among married couples expecting a child.

8. Source of Funding

None.

9. Conflict of Interest

The authors declare that there is no conflict of interest.

References

1. Pauleta JR, Pereira NM, Graça LM. Sexuality During Pregnancy. *J Sex Med.* 2010;7(1):136–42. doi:10.1111/j.1743-6109.2009.01538.x.
2. Liu HL, Hsu P, Chen KH. Sexual Activity during Pregnancy in Taiwan: A Qualitative Study. *Sex Med.* 2013;1(2):54–61.
3. Stuckey BGA. Female Sexual Function and Dysfunction in the Reproductive Years: The Influence of Endogenous and Exogenous Sex Hormones. *J Sex Med.* 2008;5(10):2282–90. doi:10.1111/j.1743-6109.2008.00992.x.
4. Basson R. human sex-response cycles. *J Sex Marital Ther.* 2001;27:33–43.
5. Erol B, Sanli O, Korkmaz D, Seyhan A, Akman T, Kadioglu A. A Cross-Sectional Study of Female Sexual Function and Dysfunction During Pregnancy. *J Sex Med.* 2007;4(5):1381–7. doi:10.1111/j.1743-6109.2007.00559.x.
6. Wan-Yim I. Childbirth among Hong Kong Chinese. In: Selin H, editor. *Childbirth across cultures: Ideas and practices of pregnancy, childbirth and post-partum.* New York: Springer; 2009. p. 71–6.
7. Xuan W. Traditional Chinese Medicine. In: Yuan CS, Bieber EJ, Bauer BA, editors. *Textbook of complimentary and alternative medicine.* 2nd edition. London: Informa Healthcare; 2006. p. 211–24.
8. Uwaputitanon W, Choobun T. Sexuality and sexual activity in pregnancy. *J Med Assoc Thai.* 2004;87(3):45–9.
9. Banerjee SK, Andersen KL, Warvadekar J, Aich P, Rawat A, Upadhyay B. How prepared are young, rural women in India to address their sexual and reproductive health needs? a cross-sectional assessment of youth in Jharkhand. *Reprod Health.* 2015;12(1):12. doi:10.1186/s12978-015-0086-8.
10. Fok WY, Chan LY, Yuen PM. Sexual behaviour and activity in Chinese women. *Acta Obstet Gynecol Scand.* 2005;84:934–8.
11. Polomeno V. Sex and Pregnancy: A Perinatal Educator's Guide. *J Perinat Educ.* 2000;9(4):15–27. doi:10.1624/105812400x87879.
12. von Sydow K. Sexuality during pregnancy and after childbirth: a metacontent analysis of 59 studies. *J Psychomatic Res.* 1999;47(1):27–9.
13. Serati M, Salvatore S, Siesto G, Cattoni E, Zanirato M, Khullar V, et al. Female Sexual Function during Pregnancy and after Childbirth. *J Sex Med.* 2010;7(8):2782–90. doi:10.1111/j.1743-6109.2010.01893.x.
14. Senkumwong N, Chaovitsaree S, Rugpao S, Chandrawongle W, Yanunto S. The changes in sexuality in Thai women during pregnancy in Thai women during pregnancy. *J Med Assoc Thai.* 2006;89(4):124–9.
15. Gokyildiz S, Beni NK. The effects of pregnancy on sexual life. *J Sex Marital Ther.* 2005;31:201–15.
16. Aslan G, Aslan D, Kızılyar A, İspahi Ç, Esen A. A prospective analysis of sexual functions during pregnancy. *Int J Impotence Res.* 2005;17(2):154–7. doi:10.1038/sj.ijir.3901288.
17. Basson R. Women's sexual dysfunction: revised and expanded definitions. *Can Med Assoc J.* 2005;172(10):1327–33. doi:10.1503/cmaj.1020174.
18. Huang YC. A comparison of sexual satisfaction and post-natal depression in the UK and Taiwan. *Int Nurs Rev.* 2006;53:197–204.
19. Lan Y, Wong DF. The role of social support in helping Chinese women with perinatal depressive symptoms cope with family conflict. *J Obstet Gynecol Neonatal Nurs.* 2008;37:556–71.
20. Naim M, Bhutto E. Sexuality during pregnancy in Pakistani women. *J Pak Med Assoc.* 2000;50:38–44.
21. Orji EO, Ogunlola IO, Fasubaa OB. Sexuality among pregnant women in South West Nigeria. *J Obstet Gynaecol.* 2002;22(2):166–8. doi:10.1080/01443610120113319.

Author biography

Chanda Rai, Senior Resident

Ananya Banerjee, Senior Resident

Manisha Meena, Senior Resident

Cite this article: Rai C, Banerjee A, Meena M. Sexual practices in pregnant women belonging to low socio-economic status at a tertiary care hospital in New Delhi, India. *Indian J Obstet Gynecol Res* 2021;8(1):90-94.